Complete Summary

GUIDELINE TITLE

Calcium x phosphate product.

BIBLIOGRAPHIC SOURCE(S)

Hawley C. Calcium x phosphate product. Nephrology 2006 Apr;11(S1):S206-8.

Hawley C. Calcium x phosphate product. Westmead NSW (Australia): CARI -Caring for Australasians with Renal Impairment; 2005 Oct. 7 p. [27 references]

GUIDELINE STATUS

This is the current release of the guideline.

COMPLETE SUMMARY CONTENT

DISCLAIMER

SCOPE METHODOLOGY - including Rating Scheme and Cost Analysis RECOMMENDATIONS EVIDENCE SUPPORTING THE RECOMMENDATIONS BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS IMPLEMENTATION OF THE GUIDELINE INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT CATEGORIES IDENTIFYING INFORMATION AND AVAILABILITY

SCOPE

DISEASE/CONDITION(S)

Chronic kidney disease

GUIDELINE CATEGORY

Management Treatment

CLINICAL SPECIALTY

Family Practice Internal Medicine Nephrology Pediatrics

INTENDED USERS

Physicians

GUIDELINE OBJECTIVE(S)

To explore the evidence to support an appropriate target range for the calcium x phosphate product in patients with renal impairment by looking at extraskeletal outcomes

TARGET POPULATION

Adults and children with chronic kidney disease

INTERVENTIONS AND PRACTICES CONSIDERED

Maintenance of appropriate serum calcium x phosphate product (considered but not recommended)

MAJOR OUTCOMES CONSIDERED

- Vascular calcification
- Cardiovascular mortality
- Mortality

METHODOLOGY

METHODS USED TO COLLECT/SELECT EVIDENCE

Searches of Electronic Databases

DESCRIPTION OF METHODS USED TO COLLECT/SELECT THE EVIDENCE

Databases searched: MeSH terms and text words for kidney dialysis were combined with MeSH terms and text words for serum calcium. This search was carried out in Medline (1966 to April Week 3 2005). The Cochrane Renal Group Trials Register was also searched for calcium and phosphate trials not indexed in Medline.

Date of searches: 30 April 2005.

NUMBER OF SOURCE DOCUMENTS

Not stated

METHODS USED TO ASSESS THE QUALITY AND STRENGTH OF THE EVIDENCE

Weighting According to a Rating Scheme (Scheme Given)

RATING SCHEME FOR THE STRENGTH OF THE EVIDENCE

Levels of Evidence

Level I: Evidence obtained from a systematic review of all relevant randomized controlled trials (RCTs)

Level II: Evidence obtained from at least one properly designed RCT

Level III: Evidence obtained from well-designed pseudo-randomized controlled trials (alternate allocation or some other method); comparative studies with concurrent controls and allocation not randomized, cohort studies, case-control studies, interrupted time series with a control group; comparative studies with historical control, two or more single arm studies, interrupted time series without a parallel control group

Level IV: Evidence obtained from case series, either post-test or pretest/post-test

METHODS USED TO ANALYZE THE EVIDENCE

Systematic Review with Evidence Tables

DESCRIPTION OF THE METHODS USED TO ANALYZE THE EVIDENCE

Not stated

METHODS USED TO FORMULATE THE RECOMMENDATIONS

Expert Consensus

DESCRIPTION OF METHODS USED TO FORMULATE THE RECOMMENDATIONS

Not stated

RATING SCHEME FOR THE STRENGTH OF THE RECOMMENDATIONS

Not applicable

COST ANALYSIS

A formal cost analysis was not performed and published cost analyses were not reviewed.

METHOD OF GUIDELINE VALIDATION

Comparison with Guidelines from Other Groups Peer Review

DESCRIPTION OF METHOD OF GUIDELINE VALIDATION

<u>Recommendations of Others</u>. Recommendations regarding calcium x phosphate product in patients with chronic kidney disease from the following groups were discussed: UK Renal Association, Canadian Society of Nephrology, European Best Practice Guidelines, and International Guidelines.

RECOMMENDATIONS

MAJOR RECOMMENDATIONS

Definitions for the levels of evidence (I–IV) can be found at the end of the "Major Recommendations" field.

Guidelines

No recommendations possible based on Level I or II evidence

Suggestions for Clinical Care

(Suggestions are based on Level III and IV evidence)

 In Stage 5 kidney disease, serum calcium (albumin-corrected) x phosphate product < 4.0 mmol/L is recommended. (Level III evidence – cohort data; strong effect; consistent finding)

Definitions:

Levels of Evidence

Level I: Evidence obtained from a systematic review of all relevant randomized controlled trials (RCTs)

Level II: Evidence obtained from at least one properly designed RCT

Level III: Evidence obtained from well-designed pseudo-randomized controlled trials (alternate allocation or some other method); comparative studies with concurrent controls and allocation not randomized, cohort studies, case-control studies, interrupted time series with a control group; comparative studies with historical control, two or more single arm studies, interrupted time series without a parallel control group

Level IV: Evidence obtained from case series, either post-test or pretest/post-test

CLINICAL ALGORITHM(S)

None provided

EVIDENCE SUPPORTING THE RECOMMENDATIONS

TYPE OF EVIDENCE SUPPORTING THE RECOMMENDATIONS

The type of supporting evidence is identified and graded for each recommendation (see "Major Recommendations").

BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS

POTENTIAL BENEFITS

Appropriate management of serum calcium x phosphate levels in patients with chronic kidney disease

POTENTIAL HARMS

Not stated

IMPLEMENTATION OF THE GUIDELINE

DESCRIPTION OF IMPLEMENTATION STRATEGY

An implementation strategy was not provided.

INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT CATEGORIES

IOM CARE NEED

Living with Illness

IOM DOMAIN

Effectiveness

IDENTIFYING INFORMATION AND AVAILABILITY

BIBLIOGRAPHIC SOURCE(S)

Hawley C. Calcium x phosphate product. Nephrology 2006 Apr;11(S1):S206-8.

Hawley C. Calcium x phosphate product. Westmead NSW (Australia): CARI - Caring for Australasians with Renal Impairment; 2005 Oct. 7 p. [27 references]

ADAPTATION

Not applicable: The guideline was not adapted from another source.

DATE RELEASED

2005 Oct

GUIDELINE DEVELOPER(S)

Caring for Australasians with Renal Impairment - Disease Specific Society

SOURCE(S) OF FUNDING

Industry-sponsored funding administered through Kidney Health Australia

GUIDELINE COMMITTEE

Not stated

COMPOSITION OF GROUP THAT AUTHORED THE GUIDELINE

Author: Carmel Hawley

FINANCIAL DISCLOSURES/CONFLICTS OF INTEREST

All guideline writers are required to fill out a declaration of conflict of interest.

GUIDELINE STATUS

This is the current release of the guideline.

GUIDELINE AVAILABILITY

Electronic copies: Available in Portable Document Format (PDF) from the <u>Caring</u> for Australasians with Renal Impairment Web site.

Print copies: Available from Caring for Australasians with Renal Impairment, Locked Bag 4001, Centre for Kidney Research, Westmead NSW, Australia 2145

AVAILABILITY OF COMPANION DOCUMENTS

The following is available:

• The CARI guidelines. A guide for writers. Caring for Australasians with Renal Impairment. 2006 May. 6 p.

Electronic copies: Available from the <u>Caring for Australasians with Renal Impairment (CARI) Web site</u>.

PATIENT RESOURCES

None available

NGC STATUS

This NGC summary was completed by ECRI Institute on June 5, 2008. The information was verified by the guideline developer on June 11, 2008.

COPYRIGHT STATEMENT

This NGC summary is based on the original guideline, which is subject to the guideline developer's copyright restrictions.

DISCLAIMER

NGC DISCLAIMER

The National Guideline Clearinghouse™ (NGC) does not develop, produce, approve, or endorse the guidelines represented on this site.

All guidelines summarized by NGC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public or private organizations, other government agencies, health care organizations or plans, and similar entities.

Guidelines represented on the NGC Web site are submitted by guideline developers, and are screened solely to determine that they meet the NGC Inclusion Criteria which may be found at http://www.guideline.gov/about/inclusion.aspx.

NGC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or clinical efficacy or effectiveness of the clinical practice guidelines and related materials represented on this site. Moreover, the views and opinions of developers or authors of guidelines represented on this site do not necessarily state or reflect those of NGC, AHRQ, or its contractor ECRI Institute, and inclusion or hosting of guidelines in NGC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding guideline content are directed to contact the guideline developer.

© 1998-2008 National Guideline Clearinghouse

Date Modified: 9/15/2008

